

Febrile, Allergic and Hypotensive Reactions (FAHR) Case Studies

2016-2022

You are free to use these examples in your teaching material or other presentations, but please do not alter the details as the copyright to this material belongs to SHOT

Misclassification of a febrile reaction results in inappropriate immediate and future management

- *A child with aplastic anaemia receiving a platelet transfusion developed a fever of 39.2°C with rigors, hypertension and tachycardia*
- *There were no allergic features*
- *He was given an antihistamine and hydrocortisone and a plan was made for prophylactic chlorphenamine before future platelet transfusions*

Unnecessary investigations for an allergic reaction

- *A male in his 30s with thalassaemia, who had a history of allergic reactions in other settings, developed rash, urticaria, facial swelling and mild hypotension after 60mL of his third unit of red cells had been transfused*
- *Transfusion was discontinued, he was given an antihistamine and hydrocortisone and his symptoms settled*
- *He was investigated with IgA levels, mast cell tryptase, repeat group and screen, direct antiglobulin test and blood cultures, none of which showed any abnormality*

Allergic reaction to an unnecessary platelet transfusion

- *A man in his 50s was transfused one adult therapeutic dose (ATD) of apheresis platelets to cover a peripherally inserted central catheter (PICC) insertion in interventional radiology*
- *He developed peri-orbital and lip swelling and a rash*
- *He was treated with intravenous (IV) hydrocortisone and chlorphenamine with resolution of his symptoms*

Future transfusion plan fails to account for reaction type

- *A woman in her 80s with transfusion-dependent anaemia required one unit of red cells following two large nose bleeds*
- *Her haemoglobin (Hb) was 68g/L with a stated target Hb of >90g/L*
- *Midway through transfusion she developed pyrexia (temperature 38°C from baseline 36.5°C), rigors and vomiting*
- *The transfusion was stopped*
- *Investigations revealed no evidence of a serological reaction*
- *On review, frequent transfusion reaction investigations had been performed previously due to similar symptoms*
- *The patient was given a plan for premedication with paracetamol, chlorphenamine, hydrocortisone and furosemide for future transfusions*

Inappropriate treatment of a febrile reaction

- *A patient in his 50s with acute myeloid leukaemia attended the haematology day unit for a routine platelet transfusion*
- *On completion he developed rigors, fever, and breathlessness. His temperature rose to 40.1°C from a baseline of 37.4°C and oxygen saturations fell to 94% on oxygen*
- *He was given IV hydrocortisone and antihistamine with little effect. He was subsequently administered 1mg adrenaline, 4.5g piperacillin with tazobactam (tazocin) (antibiotic) IV, 1g paracetamol and IV fluids*
- *His symptoms settled over the following hour, but he was admitted for observation. Blood cultures were negative and there was no rise in mast cell tryptase*

Inappropriate treatment in the presence of a potential haemolytic transfusion reaction

- *A lady in her 70s with myelodysplastic syndrome and known alloantibodies attended for a scheduled two-unit blood transfusion. The units had been crossmatched at the reference laboratory due to slight reaction on crossmatch when performed in-house.*
- *Halfway through the second unit the patient developed rigors, a rise in temperature (38.4°C from baseline 37.7°C) and elevated blood pressure (130/60 to 167/88 mmHg)*
- *The nurse stopped the transfusion and asked for medical review. The registrar prescribed 10mg antihistamine and 100mg hydrocortisone and told the nurse to continue the transfusion in 30 minutes*
- *However, the patient's symptoms worsened, and she complained of pain in her kidneys. She was given a further 100mg hydrocortisone and 1g paracetamol. Her symptoms resolved within a few hours*
- *Samples sent for serological investigation revealed no evidence of a haemolytic transfusion reaction*

Appropriate treatment

- *A man in his 20s who had suffered polytrauma received a postoperative blood transfusion*
- *After 30 minutes, routine observations revealed a temperature rise from 37.6 to 39°C*
- *He was treated with IV paracetamol and transfusion was continued. His temperature continued to reduce until returning to baseline around 12 hours post transfusion*

Febrile reaction occurring with platelets given for an erroneous result

- *A patient in her 80s was admitted for symptoms relating to a pulmonary embolism*
- *She was prescribed two units of platelets for a low platelet count (reported as $29 \times 10^9/L$)*
- *During the second unit she developed rigors, a fever of 39.2°C and an elevated heart and respiratory rate*
- *The laboratory had noted platelet clumping and had revised the report on the system however the medical team had already acted on this initial result*

Severe allergic reaction when given platelets to reverse aspirin

- *A patient in his 70s was transfused two doses of platelets in theatre*
- *He was undergoing surgery for an acute subdural haematoma and platelets were given as he was on aspirin*
- *Fifteen minutes after his second dose, the patient developed a rapid rash covering his body and hypotension unresponsive to vasopressors*
- *The patient was treated for anaphylaxis and rapid stability was achieved*

Avoid unnecessary transfusion

- *A female in her 60s was found to have a haemoglobin of 48g/L when routine blood tests were carried out at her general practice surgery*
- *She experienced severe rigors, back pain, breathlessness and felt very cold 15 minutes after being transfused a unit of red cells for symptomatic anaemia*
- *Paracetamol alone was used to treat this reaction*
- *Future management will be with intravenous (IV) iron*

Febrile reaction inappropriately treated with an antihistamine and steroid

- *A day case patient in their 60s with myelodysplasia, haemolysis and neutropenia developed a temperature rise to 39.7°C, rigors and nausea during a red cell transfusion*
- *They were treated with hydrocortisone, chlorphenamine, paracetamol, antibiotics and admitted on to a ward*
- *Future transfusion management was stated to be pre-medication with an antihistamine and steroid*
- *Although it is not clear if steroid treatment may be beneficial for the management of their haemolysis it is unlikely to prevent a further febrile-type reaction and may make infection more likely in a vulnerable, neutropenic patient*

Reducing the number of units given at each transfusion episode as a reaction prevention strategy

- *A patient in their 60s with chronic transfusion dependent anaemia received a red cell transfusion as an inpatient*
- *During the transfusion, they developed a temperature of 38°C associated with chills and rigors*
- *The rate of the transfusion was reduced and they were given paracetamol, however their symptoms reoccurred therefore the transfusion was discontinued*
- *Future management was to limit transfusion episodes to a single unit of red cells and was reported to be effective*

Use of iron to avoid the need for red cell transfusion

- *A patient in their 80s was admitted to the ambulatory care unit for a two-unit red cell transfusion for symptomatic iron deficient anaemia*
- *Chlorphenamine and ondansetron were given pre transfusion*
- *On completion of the first unit the patient developed a temperature rise of more than 2°C, rigors, nausea and was treated with paracetamol*
- *They were discharged later the same day and intravenous iron agreed as future management*
- *It is unclear what the expected benefit was of pre-transfusion chlorphenamine, however treatment with paracetamol and future management with intravenous iron are rational*
- *If intravenous iron is given prior to the development of symptoms this is likely to prevent the need for further urgent admission and red cell transfusion*

A febrile reaction appropriately treated with paracetamol

- *A patient in their 80s received a red blood cell transfusion to treat ongoing non-severe bleeding associated with a haemoglobin (Hb) of about 80g/L*
- *After 100mL had been transfused (30–60 minutes) the patient experienced rigors, an increase in respiratory rate and the temperature was noted to have risen from a baseline of 36.6°C to 38.3°C*
- *There were no other symptoms or signs*
- *The transfusion was initially slowed and then discontinued*
- *Paracetamol was prescribed and the patient's observations returned to baseline*
- *Bacterial cultures from the patient at the time of the reaction were negative*
- *No change in management was planned for any subsequent blood transfusion*

A febrile reaction to red cells. To receive iron as future management of iron deficiency anaemia

- *A patient with menorrhagia and Hb of 50g/L was transfused with red cells. After the first unit her post-transfusion observations identified a pyrexia of 39.6°C (an increase of more than 2°C from baseline) and tachycardia of 120 beats/minute*
- *She was given treatment which included paracetamol and made a complete recovery with observations returning to baseline over 1-4 hours*
- *Repeat serology was negative and future management was planned with intravenous iron and avoidance of blood transfusion*

Allergic reaction to apheresis platelets with planned transfusion of pooled platelets suspended in PAS for future management

- *A child with reversible bone marrow failure and thrombocytopenia received apheresis platelets prior to an operation*
- *Within 10 minutes of the start of the transfusion, periorbital oedema, wheezing and a fall in oxygen saturations to 92% on air occurred*
- *Oxygen therapy, hydrocortisone, chlorphenamine and salbutamol nebuliser were given with complete recovery within 1-4 hours*
- *Investigation did not identify IgA deficiency and mast cell tryptase remained within the normal range*
- *The patient had experienced previous mild reactions to apheresis platelets and so it was agreed that in future platelets suspended in PAS would be used to reduce the risk of a further allergic reaction*

A febrile reaction treated with hydrocortisone and chlorphenamine

- *An adult male with sickle cell disease attended an outpatient department to receive an exchange blood transfusion*
- *After the first unit of red cells he developed rigors. Observations revealed a temperature of 38.6°C and a rise of 2°C*
- *His blood pressure was also increased compared to pretransfusion observations but there were no respiratory signs or symptoms*
- *The transfusion was discontinued and he was given hydrocortisone and chlorphenamine*
- *He recovered in less than one hour and was subsequently admitted to the ward for antibiotics to treat a possible chest infection*
- *Repeat serology and blood culture of the patient and implicated unit were negative*

A moderate febrile reaction resulting in transfer of the patient from a community hospital to a larger hospital with an emergency department

- *An elderly male with myelodysplastic syndrome (MDS) received two units of red cells in a community hospital*
- *He was known to have anti-C, anti-Kpa and a non-specific autoantibody*
- *Following transfusion of his second unit routine observations identified a temperature rise from 36.9°C prior to transfusion to 38.7°C*
- *An ambulance was called and the patient transferred to the emergency department at a larger hospital*
- *On arrival he was given paracetamol, his temperature settled and he was discharged home*
- *Repeat serology, and culture of the patient and implicated unit revealed nil significant*

An allergic reaction to apheresis platelets

- *An elderly male with MDS and possible sepsis but no bleeding received a unit of apheresis platelets*
- *Ten minutes after starting the transfusion he developed a swollen tongue and was unable to talk*
- *His observations were stable, the transfusion was discontinued and he was given intravenous hydrocortisone*
- *The reaction resolved and a decision made that further platelet transfusion should routinely be covered with both hydrocortisone and chlorphenamine*

A severe reaction in a patient with IgA deficiency

- *An adult female received transfusion of red cells to treat a postpartum bleed on the delivery ward*
- *Within 15 minutes of the start of the transfusion she developed fever, chest tightness and throat swelling associated with a temperature rise of more than 2°C to 39.7°C, dyspnoea and visible angiodema*
- *She received paracetamol, an antihistamine, hydrocortisone and intravenous adrenaline*
- *After 4 hours her observations settled*
- *Subsequent investigation identified that she was IgA deficient with IgA antibodies and IgA deficient or washed red cells were recommended for any future transfusion*