
Paediatric Cases

FIGURES FROM THE ANNUAL SHOT REPORTS 2020-2022

You are free to use these slides in your teaching material or other presentations, but please do not alter the details as the copyright to this material belongs to SHOT




Trends in paediatric reports 2013-2022






Summary of paediatric cases by category and age 2022






Percentages of paediatric and total reports in each category in 2022



UCT=uncommon complications of transfusion; TRALI=transfusion-related acute lung injury; TAD=transfusion-associated dyspnoea; TACO=transfusion-associated circulatory overload; HTR=haemolytic transfusion reactions; FAHR=febrile, allergic and hypotensive reactions; HSE=handling and storage errors; IBCT-SRNM=incorrect blood component transfused-specific requirements not met; IBCT-WCT=IBCT-wrong component transfused




Breakdown of incorrect blood component transfused reports in 2022



IBCT-WCT=incorrect blood component transfused-wrong component transfused; IBCT-SRNM=IBCT-specific requirements not met; HSCT=haemopoietic stem cell transplant; HLA=human leucocyte antigen




Summary of FAHR reports by component type from 2013 to 2022






Comparison of proportions of adult and paediatric FAHR related to different components in 2022






Percentages of reaction types of each component for paediatric reports in 2022






Trends in paediatric reports 2012-2021






Percentages of paediatric and total reports in each category in 2021



TTI=transfusion-transmitted infection; CS=cell salvage; UCT=uncommon complications of transfusion; TRALI=transfusion-related acute lung injury; TAD=transfusion-associated dyspnoea; TACO=transfusion-associated circulatory overload; HTR=haemolytic transfusion reactions; FAHR=febrile, allergic and hypotensive reactions; HSE=handling and storage errors; PCC=prothrombin complex concentrates; IBCT-SRNM=incorrect blood component transfused-specific requirements not met; IBCT-WCT=IBCT-wrong component transfused



Summary of paediatric cases by category and age 2021



Incorrect blood component transfused (IBCT)

Avoidable transfusion

Delayed transfusion

Under or overtransfusion

- Prothrombin complex concentrate (PCC)
 - Cell salvage (CS)

Handling and storage errors (HSE)

Anti-D immunoglobulin errors (Anti-D Ig)


Febrile, allergic and hypotensive reactions (FAHR)

Haemolytic transfusion reactions (HTR)

- Transfusion-associated circulatory overload (TACO)
 - Transfusion-associated dyspnoea (TAD)
 - Uncommon complications of transfusion (UCT)




Breakdown of incorrect blood component transfused reports in 2021



Other includes incomplete testing (n=3), invalid time-expired sample (n=1), failure to provide CMV-negative (n=1) and inappropriate D-positive component (n=1) IBCT-WCT=incorrect blood component transfused-wrong component transfused; IBCT-SRNM=IBCT-specific requirements not met; CMV=cytomegalovirus




Summary of FAHR reports by component type from 2012 to 2021






Comparison of proportions of adult and paediatric FAHR related to different components in 2021






Percentages of reaction types of each component for paediatric reports in 2021






Pulmonary complications in children and neonates 2012-2021






Trends in paediatric reports from 2011-2020





Summary of paediatric cases by category and age in 2020 (n=159)



Incorrect blood component transfused (IBCT)

Under or overtransfusion

Delayed

Febrile, allergic and hypotensive reactions (FAHR)

Anti-D immunoglobulin errors (Anti-D lg)

Handling and storage errors (HSE)

Transfusion-associated circulatory overload (TACO)


Haemolytic transfusion reactions (HTR)

Transfusion-associated dyspnoea (TAD)

Uncommon complications of transfusion (UCT)




Percentages of paediatric and total reports in each category in 2020



TTI=transfusion-transmitted infection; UCT=uncommon complications of transfusion; TRALI=transfusion-related acute lung injury; TAD=transfusion-associated dyspnoea; TACO=transfusion-associated circulatory overload; HTR=haemolytic transfusion reactions; FAHR=febrile, allergic and hypotensive reactions; HSE=handling and storage errors; IBCT-SRNM=incorrect blood component transfused-specific requirements not met; IBCT-WCT=IBCT-wrong component transfused



Breakdown of incorrect blood component transfused reports in 2020 (n=44)




IBCT-WCT=incorrect blood component transfused-wrong component transfused; IBCT-SRNM=IBCT-specific requirements not met; MB=methylene blue-treated; SD=solvent-detergent treated

Note: Category 'other' includes invalid sample (n=1), K positive red cells to individual with childbearing potential (n=1), failure to provide washed platelets (n=1)




Transfusion set up for neonates and infants





Trend in paediatric FAHR reports 2011-2020




SHOT Serious Hazards of Transfusion

Paediatric FAHR reports in 2020 (n=54)

a. Proportions of adult and paediatric FAHR





b. Paediatric FAHR reports by reaction and component

